

요로결석 환자에서 요관경하 배석술 후 발생한 패혈성 속을 동반한 급성 췌장염

가톨릭 대학교 의과대학 내과학교실

박상미 · 정희선 · 김수연 · 고훈영 · 윤선애 · 김영수 · 김영옥

Acute Pancreatitis and Septic Shock Complicated after Ureteroscopic Stone Removal in a Patient with Ureter Stones

Sang Mi Park, Hee Sun Chung, Su Yoen Kim, Hun Young Ko, Sun Ae Yoon, Young Soo Kim, Young Ok Kim

The Catholic University of Korea, Medical College, Seoul, Korea

Ureteroscopy is relatively safe and one of the techniques most widely used for upper urinary tract disease. To our knowledge, acute pancreatitis complicated after ureteroscopic procedure has not reported yet. A 49-year-old male was presented with sudden onset of fever and decreased mentality just after ureteroscopic stone removal. He had received nephrothotomy 2 months ago. Urine and blood culture showed *Citrobacter freundii* and serum amylase and lipase increased to 6067 IU/L, 1270 IU/L, respectively. Abdominal CT scan demonstrated perinephric inflammatory fluid collection and its direct extension to tail portion of pancreas. With medical treatment including adequate antibiotics, urosepsis and acute pancreatitis completely recovered.