

신이식 환자에서 발생한 염증성 장질환 1례

서울시립북부노인병원 내과¹, 서울대학교병원 신장내과²

이향림¹ · 선휘경² · 구호석² · 김동기² · 오국환² · 주권욱² · 김연수²

De-novo Inflammatory Bowel Disease after Kidney Transplantation

Hyang Lim Lee¹, Hui-Kyoung Sun², Ho Seok Koo², Dongki Kim²
Kook-Hwan Oh², Kwon Wook Joo², Yon Su Kim²

Seoul Bukbu Geriatric Hospital Department of Internal Medicine¹
Seoul National University Hospital Department of Internal Medicine Division of Nephrology²

Post transplant de-novo inflammatory bowel disease (IBD) may develop despite the use of immunosuppressants including mycophenolate mofetil, a drug used for the treatment of IBD,

A 25-year old Korean man received a kidney transplantation in 1989 (at the age of 14) and has been on immunosuppression by calcort (6mg qd) and mycophenolate mofetil (500 mg bid). He was admitted to our hospital with right lower quadrant abdominal pain for 2weeks. Recently, he had bloody diarrhea. Colonoscopic examination revealed active colitis involving the terminal ileum and the ileocolic valve. Histological abnormalities were comprised of focal cryptitis, inflamed granulation tissue, lymphoid follicle and ulcer. In combine, Crohn

Key Words : 신이식, 염증성 장질환, MMF

Kidney transplantation, Inflammatory bowel disease, MMF