

A Case of Lipomyelomeningocele Complicated by Neurogenic Bladder

전남대학교 의과대학 내과학교실

서일국, 최홍상, 김민지, 나졸래, 김수희, 김창성, 최준석, 박정우, 배은희, 마성권, 김수완

A Case of Lipomyelomeningocele Complicated by Neurogenic Bladder

Il Kook Seo, Hong Sang Choi, Min Ji Kim, Jol Rae Na, Soo Hee Kim, Chang Seong Kim
Joon Seok Choi, Jeong Woo Park, Eun Hui Bae, Seong Kwon Ma, Soo Wan Kim

Department of Internal Medicine Chonnam National University Medical School

Lipomyelomeningocele is a rare congenital condition and an abnormal fat accumulation that starts below the skin and extends through an opening in the spine to the spinal cord. It forms a mass under skin and damage to adjacent spinal cord, which may have neurological consequences, such as muscle weakness or loss of sensation of the lower extremities, and bowel or bladder incontinence. A 19-year-old male visited our hospital due to urinary incontinence. The laboratory findings were as follows, blood urea nitrogen 45.3 mg/dL, serum creatinine 5.1 mg/dL and WBC >100/HPF on urine analysis. Kidney ultrasonography revealed both hydronephrosis and cortical thinning, and urodynamic study showed neurogenic bladder. Pelvic MRI showed 7 cm sized fatty mass in sacrum with tethered spinal cord indicating lipomyelomeningocele. He was transferred to neurosurgery for operation. We report a case of chronic kidney disease caused by neurogenic bladder, which was induced by lipomyelomeningocele. We suggest that evaluation for congenital abnormalities of the spine and spinal cord, such as lipomyelomeningocele, will be needed for early diagnosis and treatment in young-aged patients with urinary incontinence and decreased renal function.

Key Words: Lipomyelomeningocele, 신장손상, 방광
Lipomyelomeningocele, Kidney injury, Bladder