

S-A17

혈액투석환자에서 동맥미세석회화와 관상동맥칼슘수치의 관계

가톨릭대학교 의과대학 내과학교실

이영복, 최수진, 김영수, 윤선애, 김영옥

Arterial Micro-calcification is associated with Coronary Artery Calcium Score in Hemodialysis Patients

Yeong Bok Lee, Su Jin Choi, Young Soo Kim, Sun Ae Yoon, Young Ok Kim

Departments of Internal Medicine, The Catholic University, College of Medicine, Seoul, Korea

Background: We have reported that arterial micro-calcification (AMC) of vascular access has a negative impact on access patency and cardiovascular mortality in hemodialysis (HD) patients. Reasons behind increased cardiovascular mortality in AMC are not fully understood, but it is believed that aortic stiffness is a major contributing factor. Whereas, coronary artery calcification (CAC) is quite common in HD patients and it is known as predictor of future cardiovascular events and all cause mortality in HD patients. The aim of this study was to explore the relationship between AMC and CAC in HD patients.

Methods: One hundred HD patients who received vascular access operation were included in this study. The AMC was diagnosed by pathologic examination of arterial specimen by von Kossa stain, which was acquired during the operation. All patients underwent a multi-detector computed tomography (MDCT) imaging procedure and coronary artery calcium score (CACS) was calculated. Patients were classified into two groups, according to the CACS, as low (<100), in 40 patients, and high (≥ 100), in 60 patients. We compared CACS between the patients with and without AMC.

Results: Mean age was 65.1 ± 12.7 years and the male gender was 63 (63.0%). The incidence of AMC was 60.0% (n=60). The mean CACS was 486.8 ± 857.6 (0-5674.1), and the median value was 161.2. Patients with the positive AMC group showed a significantly higher prevalence of diabetes (85.7% vs 45.5%, $p=0.000$). Positive AMC group showed high incidence of high CACS compared to negative AMC group (76.8% vs 38.6%, $p=0.000$). By binary logistic regression, high CACS was independently associated with positive AMC (OR 7.566, 95% CI 1.771-32.322, $p=0.006$).

Conclusion: The present study suggests that AMC is closely associated with CACS in HD patients.

Key Words: 동맥미세석회화, 관상동맥수치, 혈액투석

Arterial calcification, Coronary calcium score, Hemodialysis