

## 말기신부전 환자에서 sclerostin과 말초혈관석회화 및 심혈관질환 발생의 상관관계

고려대학교 안암병원 신장내과

나영주, 김선철, 양지현, 임성윤, 김명규, 조상경, 조원용

### The Relation between Sclerostin, Peripheral Vascular Calcification, and Cardiovascular Events in ESRD Patients

Young Ju Na, Sun Chul Kim, Jihyung Yang, Sung Yoon Lim, Myung-Gyu Kim  
Sang-Kyung Jo, Wonyong Cho

Division of Nephrology, Korea University Anam Hospital

**Introduction:** Sclerostin, a Wnt antagonist produced by osteocyte, is a potent regulator of bone metabolism and a novel candidate for the bone vascular axis in chronic kidney disease patients. Although sclerostin is known as an inhibitor of vascular calcification, recent studies demonstrated conflicting results about the association between sclerostin and cardiovascular events or mortality. In this study we tested the association between sclerostin, peripheral vascular calcification, and cardiovascular events in end stage renal disease (ESRD) patient starting peritoneal dialysis.

**Methods:** In this prospective study, we included 45 ESRD patients admitted to Korea University Anam Hospital for starting peritoneal dialysis. Circulating sclerostin level was measured in all patients before the start of peritoneal dialysis. Simple vascular calcification score (SVCS) was measured using plain radiographic films of both hand and the pelvis.

**Results:** Higher sclerostin level was associated with male sex, higher left ventricle (LV) mass index, and lower LV fractional shortening in univariate analysis. ESRD patients with severe vascular calcification (SVCS  $\geq 3$ ) had significantly higher prevalence of diabetes mellitus. They had higher pulse wave velocity, alkaline phosphatase, and lower cholesterol levels, LV fractional shortening. In multivariate analysis, the presence of diabetes mellitus (OR, 44.59,  $p=0.023$ ), lower cholesterol levels (OR, 0.94,  $p=0.032$ ) were independent risk factors for severe vascular calcifications. In a multivariate Cox regression model, higher pulse wave velocity (HR 1.03,  $p=0.028$ ) was significant predictor for cardiovascular events. Although higher sclerostin group (HR 9.82,  $p=0.094$ ) and lower albumin levels (HR 0.23,  $p=0.081$ ) display a tendency of increased cardiovascular events in multivariate Cox regression analysis, there was no statistical significance.

**Conclusion:** This study showed the possible important role of sclerostin in the development of vascular calcification and cardiovascular events in PD patients. Longer term follow up with large sample size will be needed to clarify this issue.

**Key Words:** 혈관석회화, 복막투석, 심혈관계질환

Vascular calcification, Dialysis, Cardiovascular disease