

복막투석 환자에서 체내 수분 상태와 염증 및 영양 지표와의 연관성(ASKK Study)

고려대학교 내과학교실¹, 가톨릭대학교 내과학교실²
 서울대학교 내과학교실³, 인제대학교 내과학교실⁴, 전남대학교 내과학교실⁵

차진주¹ · 현영율¹ · 강영선¹ · 윤기철¹ · 한현선¹ · 최율리¹ · 유지연¹ · 김용수²
 오국환³ · 김양욱⁴ · 김영훈⁴ · 신민호⁵ · 나명윤⁵ · 김남호⁵ · 차대룡¹

Assessment of Volume Status and Its Relationship with Risk of Inflammation and Nutrition in Korean PD Patients (ASKK Study)

Jin Joo Cha¹, Young Youl Hyun¹, Young Sun Kang¹, Ki Chul Yoon¹, Hyun Sun Han¹
 Yul Ri Choi¹, Ji Yeon Yu¹, Yong Soo Kim², Kook-Hwan Oh³, Yang-Wook Kim⁴
 Yeong Hoon Kim⁴, Min ho Shin⁵, Myong yun Nah⁵, Nam-ho Kim⁵, Dae Ryong Cha¹

Department of Internal Medicine¹ Korea University

Department of Internal Medicine² the Catholic University of Korea College of Medicine

Department of Internal Medicine³ Seoul National University Hospital; Department of Internal Medicine⁴ Inje University

Department of Internal Medicine⁵ Chonnam National University

Background: As volume overload is thought to be associated with microinflammation, endothelial dysfunction, malnutrition and cardiac dysfunction, to maintain adequate volume status in advanced chronic kidney disease seems to be important to prevent worse outcomes. We evaluated (1) the markers of inflammation and malnutrition in peritoneal dialysis patients according to objective volume status, and (2) the relationship of volume overload with cardiac dysfunction.

Methods: Two hundred peritoneal dialysis patients (mean age=49.2±10.8, 32% with history of hypertension, 37.5% with history of diabetes, dialysis vintage=45.6±51.1 months) from 5 nephrology center were included in the study. We measured the hydration status using body composition monitor (BCM): total body water (TBW), overhydration (OH), intracellular water (ICW), extracellular water (ECW), lean tissue mass (LTM), adipose tissue mass (ATM), body cell mass (BCM). Demographic and laboratory data including inflammatory markers (CRP, TNF α , MCP-1, IL-6, VEGF, Fractalkine) were collected. Left ventricular hypertrophy (LVH) and calcification score were evaluated.

Results: The patients with excess ECF volume more than 2.5 L (42% of all patients) showed lower lean tissue index (p=0.021), hemoglobin (p=0.003), calcium (p=0.024), protein (0.000), albumin (p=0.000), cholesterol (p=0.024) level and higher MCP-1 (p=0.00), TNF- α (p=0.004), Glucose (p=0.014), antero-posterior calcification score (p=0.041) compared to normohydrated and mild overhydrated patients. Overhydration was positively correlated with MCP-1, VEGF, Glucose, ALP, Height, systolic blood pressure and was negatively correlated with lean tissue mass, albumin, hemoglobin level. (R²=0.557, beta=-4.112, p=0.000). ECW/ICW ratio was also positively correlated with MCP-1, height, weight discrepancy between patients weight and ideal dry body weight, and negatively with lean tissue mass index, adipose tissue index, albumin, hemoglobin (R²=0.801, beta=1.877, p=0.000). Overhydration was an independent risk factor of LVH (OR=2.573 (95% CI 1.721-3.845, p=0.000), but did not increase the risk of vascular calcification.

Conclusion: In peritoneal dialysis patients, fluid overload was associated with microinflammation, malnutrition, and possibly cardiac dysfunction. Adequate volume monitoring in dialysis patients is a cornerstone to improve outcome in peritoneal dialysis patients.

Key Words: 체내 수분 상태, 복막 투석, 체성분 계측; Volume overload, Peritoneal dialysis, Body composition monitor