

복막 투석 환자의 대사 증후군 예측에 대한 시상 복부 직경의 우월성

연세대학교 의과대학 내과학교실 신장내과

이미정 · 신동호 · 김승준 · 오형중 · 유동은 · 구향모 · 김은진 · 도화미 · 한승혁 · 최규현 · 강신욱

Sagittal Abdominal Diameter is a Stronger Predictor of Metabolic Syndrome than Anthropometric Indices in Peritoneal Dialysis Patients

Mi Jung Lee, Dong Ho Shin, Seung Jun Kim, Hyung Jung Oh, Dong Eun Yoo, Hyang Mo Koo
Eun Jin Kim, Fa Mi Doh, Seung Hyeok Han, Kyu Hun Choi, Shin-Wook Kang

Yonsei University College of Medicine Department of Internal Medicine Division of Nephrology

Background: Visceral fat is a key factor in the development of cardiovascular disease and insulin resistance. Recent studies showed the anteroposterior diameter of the abdomen, sagittal abdominal diameter (SAD) was closely associated with metabolic syndrome in general population. However, the association between SAD and metabolic syndrome has never been elucidated in patients with peritoneal dialysis (PD).

Aims: The aim of this study was to investigate whether SAD was associated with metabolic syndrome in PD patients. In addition, we evaluated whether SAD predicted metabolic syndrome better than anthropometric indices in this patients.

Methods: We undertook a cross-sectional study in 90 prevalent PD patients between February 2010 and July 2010. We assessed SAD using lateral abdominal radiograph in supine position with dialysate empty. SAD was defined as the anteroposterior diameter of the abdomen at the L4-L5 level on the radiograph. In addition, anthropometry, blood pressure, fasting glucose, insulin, and lipid profiles were measured. Metabolic syndrome was defined in accordance with the modified National Cholesterol Education Program (Adult Treatment Panel III) criteria. According to the median value of SAD (25.4 cm), patients were dichotomized into higher SAD (HS) and lower SAD (LS) groups.

Results: Compared with the LS group, the mean age, body mass index (BMI), waist circumference (WC), waist-to-hip ratio (WHR), and homeostasis model assessment – insulin resistance index were significantly higher in the HS group. In addition, metabolic syndrome was more observed in this group [20 of 45 (44.4%) vs. 7 of 45 (15.5%), $p=0.005$]. Univariate analysis showed that SAD, BMI, WC, and WHR were significantly correlated with metabolic syndrome. In a multivariate logistic regression analysis adjusted for confounding factors, SAD was an independent predictor of metabolic syndrome (odds ratio = 1.024, 95% confidence interval: 1.003–1.045, $p=0.023$). Among SAD and anthropometric indices, SAD provided the highest predictive value for metabolic syndrome [SAD; area under the receiver operating characteristic curve (AUC) 0.732, $p<0.001$, BMI; AUC 0.724, $p<0.001$, WC; AUC 0.719, $p<0.001$, WHR; AUC 0.683, $p=0.006$].

Conclusion: The present study showed SAD was more predictive for metabolic syndrome in PD patients than anthropometric indices. Therefore, SAD measured by simple abdominal radiograph could be useful to assess the risk of metabolic syndrome in patients with PD.

Key Words: 시상 복부 직경, 복막투석, 대사 증후군

Sagittal abdominal diameter, Dialysis, Metabolic syndrome