

Abstract Type : Oral

Abstract Submission No. : 1248

Loss of *Cutibacterium* is responsible for CKD-associated pruritus in patients undergoing dialysis

Jeong Geon Lee¹, Seung Seob Son², Hee Seul Jeong², Nam-Jun Cho¹, Samel Park¹, Hyo-Wook Gil¹, Hoonhee Seo³, Ho-Yeon Song³, Eun Young Lee¹

¹Department of Internal Medicine-Nephrology, Soonchunhyang University College of Medicine, Korea, Republic of

²Department of Medicine and BK21 FOUR Project, Soonchunhyang University College of Medicine, Korea, Republic of

³Department of Microbiology and Immunology, Soonchunhyang University College of Medicine, Korea, Republic of

Objectives: The pathogenesis of CKD-associated pruritus is poorly understood, although it is a serious problem worldwide. The microbiome that makes up the skin might be responsible for pruritus. Therefore, we tried to confirm the relationship between CKD-associated pruritus and skin microbiome in this study.

Methods: A total of 91 patients in Soonchunhyang University Cheonan Hospital (15 control group, 40 End-Stage Kidney Disease (ESKD) patients undergoing hemodialysis, 36 ESKD patients undergoing peritoneal dialysis) were enrolled, and swabbed in three sites; back (sebaceous area), antecubital fossa (moist area), and shin (dry area). Also, enrolled patients have done a questionnaire, which is a combination of the three methods (WI-NRS, 5-D itch scale, UP-Dial). 16S gene-based metagenomics method was applied to skin microbiome analysis.

Results: The data showed a significant microbial change in the back between the control group and the ESKD group. Here, the average composition ratio of *Cutibacterium* genus in the back was significantly lower in the ESKD group than the control group ($p < 0.05$). As a result of further analysis of the two groups of low pruritus and high pruritus among ESKD patients, here too, *Cutibacterium* was significantly lower in the high pruritus group compared to the low pruritus group ($p < 0.05$). The data showed that the skin microbiome had affected the degree of pruritus since other clinical parameters, such as age, eGFR, hemoglobin, Ca \times P, and intact parathyroid hormone didn't show significant differences.

Conclusions: In ESKD patients, back was related to the changed skin microbiome characteristics, and it was confirmed that this severity correlated with the reduction of *Cutibacterium*. This research is the first research worldwide to figure out the relationship between CKD-associated pruritus in ESKD patients and the skin microbiome.