

**Abstract Submission No.: A-0546**

**Long-term Exposure to Low Perceived Temperature in Winter Increases the Risk of Death in CKD patients**

**Ara Ko**<sup>1</sup>, Sohee Oh<sup>3</sup>, Jeonghwan Lee<sup>2</sup>, Jae-Young Byon<sup>4</sup>, Whanhee Lee<sup>5</sup>, Dong Ki Kim<sup>1</sup>, Yun Kyu Oh<sup>2</sup>, Chun Soo Lim<sup>2</sup>, Jung Pyo Lee<sup>2</sup>

<sup>1</sup>Department of Internal Medicine, Seoul National University Hospital, Korea, Republic of

<sup>2</sup>Department of Medical Research Collaborating Center, Seoul National University Boramae Medical Center, Korea, Republic of

<sup>3</sup>Department of Internal Medicine, Seoul National University Boramae Medical Center, Korea, Republic of

<sup>4</sup>Department of National Meteorological Satellite Center, Korea Meteorological Administration, Jincheon, Chungcheongbuk-do, Korea, Republic of

<sup>5</sup>Department of Data Science, Pusan National University School of Biomedical Convergence Engineering, Korea, Republic of

**Objectives :** Because of the rapid climate change, global interest about its impact on human health has been heightened. Perceived temperature (PT) is defined as a temperature that reflects the actual effect of temperature on the human body by considering the wind speed, amount of clouds, and humidity information. Due to the insufficiency of data on the health effects of long-term exposure to low temperature in chronic kidney disease (CKD) patients, we aimed to analyze the effect of PT in winter on the overall mortality among CKD patients.

**Methods :** 32,870 CKD patients in Seoul participated in a retrospective cohort at three medical centers. PT was calculated by the Staiger's equation, using the temperature of a nearby automated weather station, dew point temperature, wind velocity, and cloud amount. PT is interpolated using Kriging method and mortality was assessed by using the time-varying Cox proportional hazard model. The Cox regression model was used to assess the PT corresponding to the temperature for cold wave watch or warning

**Results :** During the 6.14 ± 3.96 years of follow-up, 6,147 (18.7%) deaths were observed Since 2000, maximal and average PT had not significantly been changed. In multivariable analysis, the hazard ratio (HR) of maximum PT was 1.091 (95% confidence interval (CI) 1.073-1.109, p-value 0.<0001), and HR of average, minimal PT were 1.042 (CI 1.018-1.065, p-value 0.0004), 1.036 (CI 1.022-1.049, p-value <0.0001) respectively. The maximum PT had the highest C-index for mortality (0.783), followed by the risk of average and minimum PT (0.780). In addition, each PT for cold wave watch and warning were -21.86°C, -25.63°C. The cold wave warning at a PT of -25.63°C indicated the risk of death as HR 1.874 (CI 1.806-1.944, p-value<0.0001) and c-index 0.807.

**Conclusions :** Exposure to lower PT during winter season could increase the risk of mortality among CKD patients.

table 1.png

		Univariable				Multivariable 1				Multivariable 2			
		HR	95% CI		C-index	HR	95% CI		C-index	HR	95% CI		C-index
Perceived temperature	average	1.034	1.017	1.050	0.508	1.049	1.028	1.071	0.794	1.042	1.018	1.065	0.780
	minimum	1.044	1.033	1.055	0.520	1.038	1.027	1.052	0.794	1.036	1.022	1.049	0.780
	maximum	1.087	1.074	1.100	0.551	1.096	1.079	1.114	0.797	1.091	1.073	1.109	0.783
Traditional temperature	average	1.052	1.031	1.073	0.511	1.086	1.059	1.114	0.795	1.075	1.046	1.105	0.780
	minimum	1.167	1.149	1.183	0.560	1.183	1.161	1.205	0.801	1.172	1.149	1.196	0.786
	maximum	1.045	1.034	1.057	0.520	1.047	1.033	1.060	0.794	1.043	1.028	1.058	0.780

table 1.png

